

Crude Oil

Crude oil is a mixture of chemicals called hydrocarbons. These are chemicals that contain hydrogen and carbon only. It made from ancient biomass, mainly plankton. Crude oil straight out of the ground is not much use, as there are too many substances in it, all with different boiling points.

Before we can use crude oil we have to separate it into its different substances. We do this by fractional distillation.

How does fractional distillation work?

- Crude oil is heated and vaporises/boils.
- Vapours rise up the column, gradually cooling and condensing.
- Hydrocarbons with different size molecules condense at different levels/temperatures
- The crude oil is separated into a series of fractions with similar numbers of carbon atoms and boiling points. These are called fractions.

As the number of carbon atoms increases:

- Molecules become larger and heavier
- Boiling point increases
- Flammability decreases (catches fire less easily)
- Viscosity increases (liquid becomes thicker)

Alkanes

Crude oil is largely made up of a family of hydrocarbons called alkanes; these contain only a single (covalent) carbon to carbon bond.

You can either represent alkanes with a molecular formula, e.g.:



Methane

Ethane


Propane

Butane


Or a displayed formula:




Methane



Ethane



Propane



Butane

[H = Hydrogen, C = Carbon, - indicates a chemical bond between atoms]

Cracking


Smaller hydrocarbons make better fuels as they are easier to ignite. However, crude oil contains a lot of longer chain hydrocarbons. To break a longer chain hydrocarbon down into a smaller one we use a process known as cracking.

Cracking

So large/long alkanes get CRACKED, which means they get broken in two.


- They are heated, turned into a vapour and passed over a hot catalyst
- Cracking produces two molecules:
 1. One shorter (useful as a fuel) alkane
 2. One alkene (used to make polymers).

Summary




Fractional Distillation Column

Below is a diagram of a fractionating column; you need to know the uses but not the names of each fraction:



Cracking

Experimental set up for cracking:



Alkenes

These hydrocarbons have at least one double bonds between the carbon atom. The general formula for alkenes is C_nH_{2n} .

Alkenes are more reactive than alkanes. They react with bromine water and make it go from orange to colourless.


Alkanes do not have a double bond so the bromine water stays orange.




Alkenes

Alkenes undergo addition reactions, this is where another element or compound is added across the double bond.

Below is an example of bromine being added across a double bond:



Bromine could be replaced in this equation with another halogen, hydrogen or water. The same type of reaction would take place, however the products formed would be different. For example, the reaction of ethene with water.



Reagent	Conditions	Product
Hydrogen	Nickel catalyst, 60°C.	Alkane
Water	Steam, high temperature, high pressure. Phosphoric acid catalyst	Alcohol
Halogen	Halogens in solution for example bromine water	Haloalkane

Alkenes

A second family of hydrocarbons is alkenes; these contain at least one double (covalent) carbon to carbon bond. The general formula for alkenes is C_nH_{2n} . Alkenes are unsaturated as there is room for 2 more hydrogens around some of the carbons. You need to know the names and structures of the first 4 alkenes. You can either represent alkenes with a molecular formula, e.g.:



Ethene

Propene

Butene

Propene

Or a displayed (structural) formula:

Name	Molecular formula	Full structural formula
Ethene	C_2H_4	$\begin{array}{c} \text{H} & \text{H} \\ & \\ \text{C} & = & \text{C} \\ & \\ \text{H} & \text{H} \end{array}$
Propene	C_3H_6	$\begin{array}{c} \text{H} & \text{H} & \text{H} \\ & & \\ \text{H}-\text{C} & - & \text{C} = & \text{C} \\ & & \\ \text{H} & & \text{H} \end{array}$
Butene	C_4H_8	$\begin{array}{c} \text{H} & \text{H} & \text{H} & \text{H} \\ & & & \\ \text{H}-\text{C} & - & \text{C} - & \text{C} = & \text{C} \\ & & & \\ \text{H} & \text{H} & & \text{H} \end{array}$
Pentene	C_5H_{10}	$\begin{array}{c} \text{H} & \text{H} & \text{H} & \text{H} & \text{H} \\ & & & & \\ \text{H}-\text{C} & - & \text{C} - & \text{C} - & \text{C} = & \text{C} \\ & & & & \\ \text{H} & \text{H} & \text{H} & & \text{H} \end{array}$

7 Organic chemistry - Knowledge organiser

Name	Group
Crude oil is a finite resource	22 Cracking needs heat and a catalyst
Crude oil is made of ancient biomass (plankton)	23 Cracking produces <u>alkenes</u> (these have a <u>double bond</u>)
Crude oil is a mixture of different sized hydrocarbons	24 Alkene + bromine water (orange) → colourless bromine water
Hydrocarbons: made of only hydrogen and carbon atoms	25 <u>Small</u> hydrocarbons are <u>very useful</u> as fuels
<u>Alkanes</u> are a type of hydrocarbon with only <u>single bonds</u>	26 <u>Alkenes</u> are used to make polymers (plastics)
General formula for alkanes: C_nH_{2n+2}	27 Alkenes are unsaturated (have at least 1 double bond)
First 4 alkanes are: Methane, ethane, propane, butane	28 First 4 alkenes: ethene, propene, butene, pentene
Alkanes are saturated (carbons have single bonds)	29 General formula for alkenes: C_nH_{2n}
<div> $\begin{array}{c} \text{H} & & \text{H} & & \text{H} \\ & & & & \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{H} \\ & & & & \\ \text{H} & & \text{H} & & \text{H} \end{array}$ </div> <p>Alkanes can be represented as: C₃H₆ or</p>	30 <div> $\begin{array}{c} \text{H} & \text{H} & \text{H} \\ & & \\ \text{H}-\text{C}-\text{C}=\text{C} \\ & & \\ \text{H} & & \text{H} \end{array}$ <p>Alkenes can be represented as: C₃H₆ or</p> </div>
0 Hydrocarbons in oil are separated into fractions	31 Alkene + oxygen → incomplete combustion (smoky flame)
1 Each fraction has a different boiling point	32 Carbon double bond reacts with: water, hydrogen and halogens
2 Distillation is: evaporating then condensing	33 Alkene + hydrogen (and catalyst) → alkane
3 Fractional distillation separates the fractions	34 Alkene + water (steam & catalyst) → alcohol
4 Fractions include: gas, petrol, diesel, kerosene	35 Alkene + halogen → a molecule with 2 halogen atoms bonded
5 Products are: solvents, lubricants, polymers, detergents	36 Alcohols have a -OH group
6 Carbon atoms can form chains and rings with other atoms	37 First 4 alcohols are: methanol, ethanol, propanol and butanol
7 Larger hydrocarbons have higher mp/bp/viscosity	38 Carboxylic acids have the functional group -COOH
8 Flammability decreases with molecular size of hydrocarbons	39 First 4 Carboxylic acids: methanoic acid, ethanoic acid, propanoic acid and butanoic acid
9 Fuel + oxygen → carbon dioxide + water	40 Alkenes can be used to make polymers e.g. poly(ethene)
10 The carbon and hydrogen are oxidised in combustion	41 Monomers are polymerised into a polymer
11 Cracking hydrocarbons breaks larger → smaller	42 The process is called ' addition polymerisation '